

Inverse volume corrections from loop quantum gravity and the primordial tensor power spectrum in slow-roll inflation

J. Grain*

*Laboratoire AstroParticule & Cosmologie,
Université Paris 7 Denis Diderot, CNRS, IN2P3
10, rue Alice Domon et Léonie Duquet,
75205 Paris cedex 13, France*

and

*Institut d'Astrophysique Spatiale,
Université Paris-Sud 11, CNRS
Bâtiments 120-121, 91405 Orsay cedex, France*

A. Barrau†

*Laboratoire de Physique Subatomique et de Cosmologie,
UJF, INPG, CNRS, IN2P3
53, avenue des Martyrs, 38026 Grenoble cedex, France*

and

*Institut des Hautes Etudes Scientifiques
35, route de Chartres, 91440 Bures-sur-Yvette, France*

A. Gorecki‡

*Laboratoire de Physique Subatomique et de Cosmologie, UJF, INPG, CNRS, IN2P3
53, avenue des Martyrs, 38026 Grenoble cedex, France*

(Dated: April 16, 2009)

Together with holonomy corrections, inverse volume terms should be taken into account when studying the primordial universe in loop quantum cosmology. We investigate how the tensor power spectrum is modified with respect to the standard general relativistic prediction by those semiclassical corrections. Depending on the values of the free parameters of the model, it is shown that the spectrum can exhibit a very large deviation from its usual shape, in particular with a very *red* slope and a strong running in the infrared limit.

PACS numbers: 04.60.Pp, 04.60.Bc, 98.80.Cq, 98.80.Qc

INTRODUCTION

A quantum theory of gravity is probably necessary to investigate situations where General Relativity (GR) breaks down. The early universe is a paradigmatic example of such a situation where the backward evolution of a classical space-time inevitably comes to an end after a finite amount of time. Among the theories willing to reconcile the Einstein gravity with quantum mechanics, Loop Quantum Gravity (LQG) is especially appealing as it is based on a nonperturbative quantization of 3-space geometry (see, *e.g.*, [1] and [2] for an introduction). Loop Quantum Cosmology (LQC) is a finite, symmetry reduced model of LQG suitable for the study of the whole Universe as a simple physical system (see, *e.g.*, [3]). On the other hand, it is well known that the inflationary scenario is currently the favored model to describe the first stages of the evolution of the Universe (see, *e.g.*, [4] for a recent review). Although still debated, it has received quite a lot of experimental confirmations, including from the WMAP 5-Years results [5], and solves most cosmological paradoxes. In this article, we consider the influence of LQC corrections to general relativity on the gravitational wave production during inflation. In this intricate framework, we assume the background to be described by the standard slow-roll inflationary scenario whereas LQC corrections are taken into account to compute the propagation of tensor modes. This approach is heuristically justified (to decouple the physical effects) and intrinsically plausible (as the LQC-driven superinflation can only be used to set the proper initial conditions to a standard inflationary stage if the horizon *and* flatness problems are both to be solved [6]). In [7, 8],

*Electronic address: julien.grain@ias.u-psud.fr

†Electronic address: aurelien.barrau@cern.ch

‡Electronic address: alexia.gorecki@lpsc.in2p3.fr

holonomy corrections (due to the fact that loop quantization is based on exponentials of the connections, rather than direct connection components) were exhaustively considered. We now focus on the other fundamental LQC correction : the inverse volume –or density-operator– (due to terms in the Hamiltonian constraint which cannot be quantized directly but only after being reexpressed as a Poisson bracket not involving an inverse). In the first section, the basic formalism is given together with the equations of motion derived in this framework. The second section deals with the definition of the power spectra and the question of initial conditions. Some analytical results are obtained in the third section. Finally, the fourth section explores the full parameter space with numerical investigations.

I. FORMALISM AND EQUATIONS OF MOTION

Loop Quantum Gravity relies on a Hamiltonian formulation of GR. The nonperturbative quantum effects associated with the LQG quantization procedure lead to effective semiclassical LQC equations. The associated hamiltonian constraint has been obtained by several different approaches [9, 10, 11, 12]. The most important step in formulating LQG is to rewrite canonical gravity in terms of Ashtekar variables, which are the densitized triad E_i^a and the Ashtekar connection A_a^i where $E_i^a = e_i^a / |\det e_i^b|$, $e_i^a e_i^b = q^{ab}$, q_{ab} is the spatial metric, and $A_a^i = \Gamma_a^i + K_a^i$ with Γ the spin connection and K the extrinsic curvature. The indices run from one to three. When written in Ashtekar variables, the matter Hamiltonian for a general space-time becomes [13]

$$\mathcal{H}_\phi = \frac{p_\phi^2}{2\sqrt{|\det E_j^c|}} + \frac{E_i^a E_i^b \partial_a \phi \partial_b \phi}{2\sqrt{|\det E_j^c|}} + \sqrt{|\det E_j^c|} V(\phi) . \quad (1)$$

The terms involving inverse expressions cannot be straightforwardly quantized and must be regularized by a dedicated procedure [14, 15]. The expressions which result from this approach are rather complicated, and, in particular, are subject to a number of quantization ambiguity parameters. In principle, the spectrum for the inverse volume can be calculated exactly in isotropic LQC but the regularization leads to some ambiguities [16]. The first important scale is set by $\ell_i = \sqrt{\gamma} \ell_{Pl}$, where γ is the Barbero-Immirzi parameter. It can be understood as the length above which space-time is roughly continuous and the inverse spectrum can be described by a continuous function. The second relevant scale is $\ell_* = \ell_i \sqrt{j}/3$, where j , which takes half integer values, is one of the ambiguity parameters. Above this scale the eigenvalues of the inverse operator follow the classical values, while they radically differ below ℓ_* .

Loop Quantum Gravity introduces strong modifications to the dynamical equations in the semiclassical regime (*i.e.* when the scale factor a is such that $\ell_{Pl} < a < \ell_*$). They come from the density operator [17]

$$d_j(a) = \frac{D(q)}{a^3},$$

with $q = (a/\ell_*)^2$. For the semiclassical universe, the quantum correction factor is given by [18]

$$D(q) = q^{\frac{3}{2}} \left\{ \frac{3}{2m} \left[\frac{1}{m+2} \left(|q+1|^{m+2} - |q-1|^{m+2} \right) - \frac{q}{m+1} \left(|q+1|^{m+1} - \text{sgn}(q-1) |q-1|^{m+1} \right) \right] \right\}^{\frac{3}{2-2m}}, \quad (2)$$

m being an ambiguity parameter satisfying $0 < m < 1$. The cosmological dynamics with a field χ is then governed by the following set of differential equations [19] :

$$\begin{aligned} \ddot{\chi} + \left(3H - \frac{\dot{D}}{D} \right) \dot{\chi} + D \frac{dV}{d\chi} &= 0 \\ H^2 - \frac{8\pi G}{3} \left[\frac{\dot{\chi}^2}{2D} + V(\chi) \right] &= 0 \\ \frac{\ddot{a}}{a} + \frac{8\pi G}{3} \left[\frac{\dot{\chi}^2}{D} \left(1 - \frac{\dot{D}}{4HD} \right) - V(\chi) \right] &= 0, \end{aligned} \quad (3)$$

where $H = \dot{a}/a$ is the Hubble constant and a *dot* means differentiation according to the cosmic time. When $a \geq \ell_*$, the Universe enters its classical regime and, in the limit $a \gg \ell_*$, leading to $D(q) \sim 1$, the usual Klein-Gordon equation is recovered for the inflaton field. In the semiclassical regime, $a < \ell_*$, it has, however, been shown that spectacular

modifications to the standard dynamics can be expected. For a scalar field driven dynamics, it seems that the field can naturally be excited up its self-interaction potential, setting the initial conditions for slow-roll inflation (see, *e.g.*, [6, 20, 21, 22, 23, 24, 25]). This is a very appealing feature of LQC as the requirement $\phi_i \geq 3M_{Pl}$ (for simple ϕ^2 inflationary potential), imposed by observations, is rather difficult to set in the standard framework. Furthermore, a period of superinflation ($\dot{H} > 0$) is expected to generically occur (see, *e.g.*, [26, 27, 28]), irrespectively of the detailed shape of the potential.

Following the notation of [25, 28], the equation of motion for tensor modes with quantum corrections coming from the density operator is given by

$$\ddot{h} + \left(3H - \frac{\dot{S}}{S}\right) \dot{h} - \frac{S^2}{a^2} \nabla^2 h = 0, \quad (4)$$

We note that such corrections are alternatively denoted by α in [29]. Because of quantum corrections encoded in the S -term, the standard transformation from cosmic time t to conformal time η ($d\eta = dt/(a(t))$) with the usual field redefinition $\Phi = ah$, does *not* lead to a Schrödinger equation (see, *e.g.*, Eq. (26) in [19]). In particular, there is an additional (anti)friction term given by $-\dot{S}/S$. To recast the aforementioned differential equation into a Schrödinger-like equation, we switch from cosmic time to conformal time and redefine the field according to

$$\Phi(\eta, \vec{x}) = \frac{a(\eta)}{\sqrt{S(\eta)}} h(\eta, \vec{x}).$$

Finally, by decomposing the field Φ over its spatial Fourier modes, one obtains

$$\left[\frac{d^2}{d\eta^2} + S^2(\eta)k^2 - V_1(\eta) \right] \phi_k(\eta) = 0, \quad (5)$$

with a potential term given by

$$V_1(\eta) = \frac{a''}{a} - \frac{a'}{a} \frac{S'}{S} + \frac{3}{4} \left(\frac{S'}{S} \right)^2 - \frac{1}{2} \frac{S''}{S}. \quad (6)$$

The *prime* should also be understood as a differentiation according to the conformal time. In the classical regime ($S \sim 1$), the potential term becomes $V \approx a''/a$, which is the usual GR expression.

In addition to density-operator corrections, gravitational waves propagating in a FLRW background receive quantum corrections from *holonomies* [29]. The influence of these LQC corrections has been studied in [7, 8].

The values of ambiguity parameter n , which depends on the scheme adopted to quantize holonomies, range between $-1/2$ and 0 , though it was recently shown that $n = -1/2$ is favored [34]. Moreover, if $n > -1/2$, the holonomy corrections may become a major contribution to the effective mass of the gravitons at the end of inflation [8], leading to a rather intricate picture. In the case $n = -1/2$, the potential term reads:

$$V_2(\eta) = \frac{a''}{a} - 2\sqrt{\pi}\gamma \frac{\gamma^2}{M_{Pl}^2} \left[\frac{3}{2\gamma^2 L^2(\eta)} \left(1 - \sqrt{1 - 4\frac{H^2(\eta)}{\gamma^2 L^2(\eta)}} \right) \right]^2 a^2, \quad (7)$$

where L is the comoving size of a given patch.

We will resist the temptation to combine V_1 and V_2 into a single equation which would describe both holonomy and inverse-volume corrections to the propagation of gravitational waves as our approach is to decouple, as much as possible, the different physical effects. This is, however, the next logical step in this study. Whatever the class of quantum correction considered, the equation of motion for primordial gravitons can be written as a Schrödinger-like equation:

$$\left[\frac{\partial^2}{\partial \eta^2} + E_k(\eta) - V(\eta) \right] f_k(\eta) = 0. \quad (8)$$

The difference $[E_k(\eta) - V(\eta)]$ can be seen as the effective squared-frequency $\omega_k^2(\eta)$. The correspondence between the different terms is summarized in Table I. If $E_k(\eta) > V(\eta)$, the solution oscillates whereas it becomes a coherent sum

	Density-operator corrections	Holonomy corrections
f_k	$\phi_k \equiv ah/\sqrt{S}$	$\psi_k \equiv ah$
$E_k(\eta)$	$S^2(\eta)k^2$	k^2
$V(\eta)$	$V_1(\eta)$	$V_2(\eta)$

TABLE I: Correspondence between the energylike and potential-like term.

of an evanescent and an exponentially increasing wave if $E_k(\eta) < V(\eta)$. Amplification of quantum fluctuations then arises when $E_k(\eta) < V(\eta)$, *i.e.* when ω_k^2 is negatively valued.

From now on, we switch to our main working hypothesis: as in [8], the background is supposed to be classical (*i.e.* described by the usual slow-roll inflationary picture) whereas the mode propagation is corrected by the inverse-volume LQC term. This makes sense as a phase of standard inflation is anyway mandatory after the superinflation regime. Furthermore, this allows us to understand in details the physical origin of the observed features. During slow-roll inflation, the scale factor is given by

$$a(\eta) = \ell_0 |\eta|^{-1-\epsilon},$$

with ϵ the first parameter of the slow-roll expansion. The energy density is assumed to be related to the Hubble parameter via the standard Friedmann equation. The S operator is given by [29]

$$S(q > 1) \approx 1 + \lambda q^{-\kappa/2}, \quad (9)$$

with λ and κ two positive constants, not well constrained in homogeneous models ($\kappa = 2n$ with the notation of [29]). Setting $\ell_* \sim \ell_{Pl}$ to remain consistent with our hypothesis of a full "standard" (*i.e.* classical) evolution of the background, one obtains the following energy and potential terms for density operator corrections:

$$E_k(\eta) = \left[1 + 2\lambda \left(\frac{\ell_{Pl}}{\ell_0} \right)^\kappa |\eta|^{\kappa(1+\epsilon)} \right] k^2, \quad (10)$$

$$V(\eta) = \frac{2+3\epsilon}{\eta^2} + \lambda\kappa(1+2\epsilon) \left(\frac{\ell_{Pl}}{\ell_0} \right)^\kappa |\eta|^{\kappa(1+\epsilon)-2}, \quad (11)$$

and, for holonomy corrections:

$$E_k(\eta) = k^2, \quad (12)$$

$$V(\eta) = \frac{2+3\epsilon}{\eta^2} - 2\sqrt{\pi}\gamma^3(1+4\epsilon) \left(\frac{\ell_{Pl}}{\ell_0} \right)^2 |\eta|^{2\epsilon-2}. \quad (13)$$

II. ANALYTIC RESULTS

The general equation of motion is far too complicated to be analytically solved in the general case. However, for some particular values of the parameters, the spectrum can be computed, at least in the infrared (IR) or ultraviolet (UV) limits. Those calculations are both useful by themselves and convenient to check the numerical results obtained in the next section.

Throughout all this article, the convention of [33] is used for the normalization of initial states. The particle interpretation of the considered quantum field theory imposes the Wronskian W of the mode functions to be equal to $-i$. However, because we are working with rescaled quantities, the mode functions have to be normalized so that:

$$\begin{aligned} W_\eta(\phi^\dagger, \phi) &\equiv \phi \partial_\eta \phi^\dagger - \phi^\dagger \partial_\eta \phi \\ &= -\frac{16i\pi}{M_{Pl}}. \end{aligned} \quad (14)$$

The power spectrum then reads:

$$\mathcal{P}_T(k) = \frac{2k^3}{\pi^2} \left| \frac{\sqrt{S}\phi_k}{a} \right|_{k/aH \rightarrow 0}^2. \quad (15)$$

As $S \sim 1$ at the end of inflation, the power spectrum can be safely approximated by

$$\mathcal{P}_T(k) = \frac{2k^3}{\pi^2} \left| \frac{\phi_k}{a} \right|_{k/aH \rightarrow 0}^2. \quad (16)$$

A. $\kappa = 1$ and $\epsilon = 0$: UV limit

When $\kappa = 1$ and $\epsilon = 0$, the UV limit of the power spectrum can be analytically computed. The equation of motion can be rewritten as

$$\frac{d^2\phi}{d\eta^2} + \left[(1 + 2Z|\eta|)k^2 - \frac{2}{\eta^2} - \frac{Z}{|\eta|} \right] \phi = 0, \quad (17)$$

with $Z \equiv \lambda(\ell_{\text{Pl}}/\ell_0)^\kappa = \lambda(\ell_{\text{Pl}}/\ell_0)$ in this case.

It is useful to define two regions: region I is such that the potential term can be neglected and region II is such that the time-dependent term in the energy can be neglected. Obviously, region I corresponds to $\eta \ll \eta_c$ where η_c is the potential crossing time defined by $E(\eta_c) = V(\eta_c)$. Region II corresponds to $-Z^{-1} \ll \eta \leq 0$. In region I, the mode functions are given by a linear combination (LC) of Airy functions:

$$\phi_I(\eta) \sim A_I \mathbf{Ai}(x) + B_I \mathbf{Bi}(x), \quad (18)$$

with

$$x = -k^{2/3}(2Z)^{-2/3}(1 + 2Z|\eta|). \quad (19)$$

The two coefficients are determined by the standard Wronskian condition

$$(2Zk^2)^{1/3} \frac{2i\text{Im}[A_I^\dagger B_I]}{\pi} = -i \frac{16\pi}{M_{\text{Pl}}^2}, \quad (20)$$

leading to the natural choice

$$A_I = \frac{2\pi\sqrt{2}}{(k\sqrt{2Z})^{1/3} M_{\text{Pl}}}, \quad (21)$$

$$B_I = -i \frac{2\pi\sqrt{2}}{(k\sqrt{2Z})^{1/3} M_{\text{Pl}}}. \quad (22)$$

In region II, the mode functions are given by a LC of Coulomb wavefunctions:

$$\phi_{II}(\eta) \sim A_{II} F_1(Z/2k, k|\eta|) + B_{II} G_1(Z/2k, k|\eta|). \quad (23)$$

In the UV regime (*i.e.* $k \rightarrow \infty$), the potential crossing time is roughly given by $\eta_c \propto -1/k$ and region I extends to values of η higher than $-1/Z$. As a consequence, the regions overlap and the matching can be done for conformal times such that

$$k^{-1} \ll |\eta| \ll Z^{-1}.$$

This region corresponds to the time when the density-operator is close to unity until horizon crossing. In this overlapping region, the Coulomb functions take the following approximate form:

$$\phi_{II} \sim [A_{II} \sin(-k\eta + \varphi_{II}) + B_{II} \cos(-k\eta + \varphi_{II})], \quad (24)$$

with

$$\varphi_{II} = -\frac{\pi}{2} - \frac{\gamma_e Z}{2k} + \arctan\left(\frac{Z}{4k}\right),$$

where γ_e is the Euler's constant and $\varphi_{II} \sim -\pi/2$ in the UV limit. On the other hand, still in the UV regime, the Airy functions can be approximated by sine and cosine functions:

$$\lim_{x \rightarrow -\infty} \mathbf{Ai}(x) = \frac{1}{\sqrt{\pi\sqrt{|x|}}} \cos\left(\frac{2}{3}|x|^{3/2} - \frac{\pi}{4}\right), \quad (25)$$

$$\lim_{x \rightarrow -\infty} \mathbf{Bi}(x) = -\frac{1}{\sqrt{\pi\sqrt{|x|}}} \sin\left(\frac{2}{3}|x|^{3/2} - \frac{\pi}{4}\right). \quad (26)$$

Taking the limit $x \rightarrow \infty$ (as $k \rightarrow \infty$ and η is in the overlapping region) and performing a Taylor expansion in $Z|\eta|$, this leads to

$$\phi_I \sim \frac{1}{\sqrt{\pi}} \left(\frac{2Z}{k}\right)^{1/6} \{[A_I \cos(\varphi_I) - B_I \sin(\varphi_I)] \cos(-k\eta) + [A_I \sin(\varphi_I) + B_I \cos(\varphi_I)] \sin(-k\eta)\}, \quad (27)$$

with $\varphi_I = k/3Z - \pi/4$.

After matching the solutions, one easily obtains the coefficients in region II:

$$A_{II} = \frac{2i\sqrt{2\pi}}{M_{\text{Pl}}\sqrt{k}} e^{-i\varphi}, \quad (28)$$

$$B_{II} = iA_{II}^\dagger, \quad (29)$$

where the UV limit of φ_{II} has been used. The power spectrum is derived by taking the asymptotic limit for small arguments of Eq. (23). This can be performed by noticing that in the UV regime, the Coulomb wave functions are well approximated by Bessel functions. This leads to

$$\mathcal{P}_T(k) = \left(\frac{\ell_{\text{Pl}}}{\ell_0}\right)^2 \left(\frac{8\Gamma(3/2)}{\pi}\right)^2, \quad (30)$$

which coincides with the power spectrum in GR.

B. $\kappa(1+\epsilon) = 2$: full spectrum and IR & UV limits

In this particular case, it is possible to calculate the full power spectrum and to derive explicit IR and UV limits. Some algebra is, however, necessary.

The equation to be solved is the following:

$$\frac{d^2\phi_k}{d\eta^2} + \left\{ \left[1 + 2\lambda \left(\frac{\ell_{\text{Pl}}}{\ell_0}\right)^\kappa |\eta|^{\kappa(1+\epsilon)} \right] k^2 - \frac{2+3\epsilon}{\eta^2} - \lambda\kappa(1+2\epsilon) \left(\frac{\ell_{\text{Pl}}}{\ell_0}\right)^\kappa |\eta|^{\kappa(1+\epsilon)-2} \right\} \phi_k = 0. \quad (31)$$

With

$$Z = \lambda \left(\frac{\ell_{\text{Pl}}}{\ell_0}\right)^\kappa, \quad (32)$$

$$K^2 = k^2 - \lambda\kappa(1+2\epsilon) \left(\frac{\ell_{\text{Pl}}}{\ell_0}\right)^\kappa, \quad (33)$$

this can be written as

$$\frac{d^2\phi_k}{d\eta^2} + \left[K^2 + 2Zk^2|\eta|^2 - \frac{2+3\epsilon}{\eta^2} \right] \phi_k = 0. \quad (34)$$

To solve this equation, we will express it as a *General Confluent Equation* whose general form is given by

$$\begin{aligned} \frac{d^2 w}{dz^2} + \underbrace{\left[\frac{2A}{z} + 2f' + \frac{bh'}{h} - h' - \frac{h''}{h'} \right]}_{\Lambda_1} \frac{dw}{dz} \\ + \underbrace{\left[\left(\frac{bh'}{h} - h' - \frac{h''}{h'} \right) \left(\frac{A}{z} + f' \right) + \frac{A(A-1)}{z^2} + \frac{2Af'}{z} + f'' + f'^2 - \frac{ah'^2}{h} \right]}_{\Lambda_2} w = 0, \end{aligned} \quad (35)$$

and whose solution is

$$w(z) = z^{-A} e^{-f(z)} \left[C_1 M(a, b, h(z)) + C_2 U(a, b, h(z)) \right], \quad (36)$$

M , U being Kummer functions and C_i being integration constants.

To write Eq. (34) as Eq. (36), Λ_1 must vanish. We define

$$h(z) = \beta z^\alpha \quad (37)$$

$$f(z) = \frac{h(z)}{2}, \quad (38)$$

where α and β are constants. The requirement $\Lambda_1 = 0$ reads as

$$2A + \alpha b - \alpha + 1 = 0. \quad (39)$$

Λ_2 must then be equaled to the $(E_k(\eta) - V(\eta))$ term in Eq. (36). With our choice for h and f , Λ_2 reads:

$$\frac{A(A-1+b\alpha-\alpha+1)}{z^2} - \left(\alpha^2 \beta \left(a - \frac{b}{2} \right) - \frac{\alpha^2 \beta^2 z^\alpha}{4} \right) z^{\alpha-2}. \quad (40)$$

Taking $\alpha = 2$, this can be written as

$$\Lambda_2 = \frac{A(A+2b-2)}{z^2} - 4\beta \left(a - \frac{b}{2} \right) - \beta^2 z^2, \quad (41)$$

which equals $E_k(\eta) - V(\eta)$ if $z = |\eta|$. Identifying the other terms leads to

$$z = |\eta|, \quad (42)$$

$$\alpha = 2, \quad (43)$$

$$\beta^2 = -2Zk^2 \quad (44)$$

$$4\beta \left(a - \frac{b}{2} \right) = -K^2, \quad (45)$$

$$A(A+2b-2) = -2-3\epsilon, \quad (46)$$

$$2A+2b-1 = 0. \quad (47)$$

This can be easily solved to obtain

$$A = -\frac{1}{2} \pm \frac{3}{2} \sqrt{1 + \frac{12}{9}\epsilon} \quad (48)$$

$$b = 1 \mp \frac{3}{2} \sqrt{1 + \frac{12}{9}\epsilon} \quad (49)$$

$$\beta = \pm ik\sqrt{2Z} \quad (50)$$

$$a = \frac{1}{2} \mp \frac{3}{4} \sqrt{1 + \frac{12}{9}\epsilon} \mp \frac{K^2}{4ik\sqrt{2Z}}. \quad (51)$$

To determine the signs, we require the solution to converge to the usual GR solution when the LQC terms are vanishing. This means:

$$\lim_{Z \rightarrow 0} \phi_k = \sqrt{z} [D_1 J_\nu(kz) + D_2 Y_\nu(kz)], \quad (52)$$

with

$$\nu = \frac{3}{2} \sqrt{1 + \frac{12}{9}\epsilon} = \mp(b-1). \quad (53)$$

When $Z \rightarrow 0$, one obtains

$$a \equiv \mp \frac{k}{4i\sqrt{2Z}}, \quad (54)$$

which tends to $\mp i\infty$. In the limit $|a| \rightarrow \infty$, Kummer functions can be rewritten with Bessel functions:

$$\lim_{Z \rightarrow 0} [M(a, b, h(z))] = \Gamma(b) \left(\frac{kz}{2} \right)^{A+1/2} J_{\mp\nu}(kz), \quad (55)$$

and b can be reexpressed as a function of either A or ν . When plugged into the general solution with Kummer functions this leads to

$$\lim_{Z \rightarrow 0} \phi_k = \sqrt{z} [E_1 J_{\mp\nu}(kz) + E_2 Y_{\mp\nu}(kz)] [1 + \mathcal{O}(z^2)], \quad (56)$$

where E_i are constants expressed with Γ functions and the C_i coefficients. To ensure the equality with Eq. (52), the (+) sign must be chosen for b and the (−) sign for A (the sign of β is not relevant). The solution to Eq. (34) is therefore

$$\phi_k(\eta) = \frac{\exp\left(-ik\sqrt{\frac{Z}{2}}\eta^2\right)}{|\eta|^{-\frac{1}{2}-\frac{3}{2}\sqrt{1+\frac{12}{9}\epsilon}}} \left[C_1 M\left(\frac{1}{2} + \frac{3}{4}\sqrt{1+\frac{12}{9}\epsilon} + \frac{iK^2}{4k\sqrt{2Z}}, 1 + \frac{3}{2}\sqrt{1+\frac{12}{9}\epsilon}, ik\sqrt{2Z}\eta^2\right) \right. \quad (57)$$

$$\left. + C_2 U\left(\frac{1}{2} + \frac{3}{4}\sqrt{1+\frac{12}{9}\epsilon} + \frac{iK^2}{4k\sqrt{2Z}}, 1 + \frac{3}{2}\sqrt{1+\frac{12}{9}\epsilon}, ik\sqrt{2Z}\eta^2\right) \right]. \quad (58)$$

After performing a first-order Taylor expansion in ϵ , the parameters are given by

$$A = -2 - \epsilon, \quad (59)$$

$$b = \frac{5}{2} + \epsilon, \quad (60)$$

$$a = \frac{5}{4} + \frac{\epsilon}{2} + \frac{iK^2}{4k\sqrt{2Z}}. \quad (61)$$

To explicitly derive the power spectrum, the solution is rewritten as

$$\phi_k(\eta) = \frac{1}{x^A} \exp\left(-ik\sqrt{\frac{Z}{2}}x^2\right) [C_1 M(a, b, ik\sqrt{2Z}x^2) + C_2 U(a, b, ik\sqrt{2Z}x^2)], \quad (62)$$

where $x = |\eta|$. The integration constants are determined by the Wronskian condition. To compute the Wronskian, we will set $h = k\sqrt{2Z}x^2$ and focus on the remote past. The solution is

$$\phi_k(h) = \left(k\sqrt{2Z}\right)^{A/2} h^{-A/2} e^{-i\frac{h}{2}} [C_1 M(a, b, ih) + C_2 U(a, b, ih)], \quad (63)$$

$$\simeq \left(k\sqrt{2Z}\right)^{A/2} \frac{(i)^{\text{Re}[-a]}}{h^{1/4}} \left\{ \lambda \exp\left(i\frac{h}{2} + i\text{Im}[a] \ln(h)\right) + \mu \exp\left(-i\frac{h}{2} - i\text{Im}[a] \ln(h)\right) \right\}, \quad (64)$$

with

$$\lambda = e^{-\pi\text{Im}[a]/2} C_1 \frac{\Gamma(b)}{\Gamma(a)} \quad (65)$$

$$\mu = e^{\pi\text{Im}[a]/2} \left[C_2 + C_1 e^{i\pi a} \frac{\Gamma(b)}{\Gamma(a-b)} \right]. \quad (66)$$

In the remote pas, the Wronskian can be written as

$$\sqrt{h}W_h \left\{ \phi_k^\dagger, \phi_k \right\} \approx -i \left(k\sqrt{2Z} \right)^A \left[|\lambda|^2 - |\mu|^2 \right]. \quad (67)$$

And the Wronskian condition

$$\sqrt{h}W_h \left\{ \phi_k^\dagger, \phi_k \right\} = \frac{8i\pi}{M_{\text{Pl}}^2 \sqrt{k\sqrt{2Z}}}, \quad (68)$$

leads to

$$|\lambda|^2 - |\mu|^2 = -\frac{8\pi}{M_{\text{Pl}}^2 \left(k\sqrt{2Z} \right)^{A+1/2}}. \quad (69)$$

Setting $\lambda = 0$, this means that

$$C_1 = 0 \quad (70)$$

$$C_2 = \frac{2\sqrt{2\pi}}{M_{\text{Pl}}} \left(k\sqrt{2Z} \right)^{-A/2-1/4} e^{-\pi \text{Im}[a]/2}. \quad (71)$$

To obtain the power spectrum, the limit $\eta \rightarrow 0$ is taken in Eq (62):

$$\begin{aligned} \phi_k(\eta \rightarrow 0) &\simeq C_2 |\eta|^{-A} \frac{\Gamma(b-1)}{\Gamma(a)} \left(ik\sqrt{2Z}\eta^2 \right)^{1-b} \\ &\simeq C_2 \frac{\Gamma(\frac{3}{2} + \epsilon)}{\Gamma(\frac{5}{4} + \frac{\epsilon}{2} + \frac{iK^2}{4k\sqrt{2Z}})} \left(ik\sqrt{2Z} \right)^{-\frac{3}{2}-\epsilon} |\eta|^{-1-\epsilon} \\ &\simeq C_2 \frac{\Gamma(\frac{3}{2} + \epsilon)}{\Gamma(\frac{5}{4} + \frac{\epsilon}{2} + \frac{iK^2}{4k\sqrt{2Z}})} \left(ik\sqrt{2Z} \right)^{-\frac{3}{2}-\epsilon} \frac{a(\eta)}{\ell_0}. \end{aligned} \quad (72)$$

The spectrum is finally given by

$$\mathcal{P}_{\text{T}}(k) = \left(\frac{\ell_{\text{Pl}}}{\ell_0} \right)^2 \frac{16}{\pi} |\Gamma(3/2 + \epsilon)|^2 (2Z)^{-\epsilon/2-3/4} \frac{k^{3/2-\epsilon} \exp\left(-\frac{\pi K^2}{4k\sqrt{2Z}}\right)}{\left| \Gamma\left(\frac{5}{4} + \frac{\epsilon}{2} + \frac{iK^2}{4k\sqrt{2Z}}\right) \right|^2}. \quad (73)$$

This establishes the full tensor power spectrum. In the IR and UV limits, $\text{Im}(a) \rightarrow \pm\infty$, so

$$\left| \Gamma\left(\frac{5}{4} + \frac{\epsilon}{2} + \frac{iK^2}{4k\sqrt{2Z}}\right) \right|^2 \simeq 2\pi \exp\left(-\left| \frac{\pi K^2}{4k\sqrt{2Z}} \right| \right) \left| \frac{K^2}{4k\sqrt{2Z}} \right|^{3/2+\epsilon}. \quad (74)$$

After performing the Taylor expansion in ϵ and taking the *IR* and *UV* values of K^2 , one obtains

$$\mathcal{P}_{\text{T}}^{(IR)} = \left(\frac{\ell_{\text{Pl}}}{\ell_0} \right)^2 \left(\frac{2^{\frac{3}{2}} \Gamma(3/2 + \epsilon)}{\pi} \right)^2 [2Z(1 + \epsilon)]^{-\frac{3}{2}-\epsilon} k^3 \exp\left(\frac{\pi\sqrt{2Z}(1 + \epsilon)}{2k}\right) \quad (75)$$

$$\mathcal{P}_{\text{T}}^{(UV)} = \left(\frac{\ell_{\text{Pl}}}{\ell_0} \right)^2 \left(\frac{2^{3+\epsilon} \Gamma(3/2 + \epsilon)}{\pi} \right)^2 k^{-2\epsilon} \left[1 + (3 + 5\epsilon) \frac{Z}{k^2} + \mathcal{O}(k^{-4}) \right]. \quad (76)$$

It can easily be seen from those equations that important modifications to the usual GR picture are expected in the IR regime whereas the UV behavior is equivalent to the GR one up to $\mathcal{O}(k^{-2})$. In the IR range, one can both notice a very red spectrum and a very strong running of the index.

The tilt of the spectrum is given by

$$n_{\text{T}}^{(IR)} = 3 - \frac{\pi\sqrt{2Z}(1+\epsilon)}{2k} \quad (77)$$

$$n_{\text{T}}^{(UV)} = -2\epsilon - (3+5\epsilon)\frac{2Z}{k^2}, \quad (78)$$

and the running by

$$\alpha_{\text{T}}^{(IR)} = \frac{\pi\sqrt{2Z}(1+\epsilon)}{2k} \quad (79)$$

$$\alpha_{\text{T}}^{(UV)} = (3+5\epsilon)\frac{4Z}{k^2}. \quad (80)$$

III. NUMERICAL RESULTS

To explore the full parameter space beyond the particular case $\kappa(1+\epsilon) = 2$, there is probably no other way to go than to perform a full numerical investigation of the problem. To this aim, a dedicated code was developed. Initial and final conditions, defined respectively in the remote past and at the end of inflation, are however required to perform such a computation.

A. Boundary conditions

To define the initial states, we first perform the following transformation:

$$|\eta| = e^x, \quad \phi_k = u(x)e^{x/2}. \quad (81)$$

The differential equation (8) with E and V given by Eq. (10) and Eq. (11) now reads:

$$\frac{d^2u}{dx^2} + \left[\left(1 + 2\lambda \left(\frac{\ell_{\text{Pl}}}{\ell_0} \right)^\kappa e^{\kappa(1+\epsilon)x} \right) e^{2x} k^2 - \lambda\kappa(1+2\epsilon) \left(\frac{\ell_{\text{Pl}}}{\ell_0} \right)^\kappa e^{\kappa(1+\epsilon)x} - \frac{9}{4} - 3\epsilon \right] u = 0. \quad (82)$$

In the remote past, the new variable x tends to infinity and the squared frequency in the above Schrödinger-like equation becomes dominated by the term proportional to $e^{\kappa(1+\epsilon)x+2x}k^2$. With this approximation and using $y = [\kappa(1+\epsilon)x + 2x]/2$, it becomes a Bessel equation. For $\eta \rightarrow -\infty$, the mode functions can therefore be written as a LC of Hankel functions of order 0:

$$\phi_k(\eta \rightarrow -\infty) = \sqrt{-\eta} \left[A_k H_0(z_\eta) + B_k H_0^\dagger(z_\eta) \right], \quad (83)$$

with

$$z_\eta = \frac{2\sqrt{2\lambda}}{\kappa(1+\epsilon)+2} \left(\frac{\ell_{\text{Pl}}}{\ell_0} \right)^{\frac{\kappa}{2}} k |\eta|^{[\kappa(1+\epsilon)+2]/2}. \quad (84)$$

The amplitudes of the mode functions are determined by requiring the appropriate Wronskian condition as defined in Eq. (14). Using

$$W_z(H_0^\dagger(z), H_0(z)) = \frac{4i}{\pi z},$$

it can be shown that the Wronskian is given by

$$W_\eta(\phi^\dagger, \phi) = -\frac{2i}{\pi} [\kappa(1+\epsilon) + 2] \left[|A_k|^2 - |B_k|^2 \right]. \quad (85)$$

The natural choice

$$A_k = \frac{2\pi\sqrt{2}}{[\kappa(1+\epsilon) + 2] M_{\text{Pl}}}, \quad (86)$$

$$B_k = 0 \quad (87)$$




FIG. 1: Primordial tensor power spectrum as a function of the wavenumber. The solid line is the analytical calculation whereas filled circles are the numerical results.

will therefore be made to fulfill the Wronskian condition.

At the very end of inflation, one can notice that whatever the LQC correction considered, it becomes subdominant when compared with the GR term. As a consequence, the equation of motion becomes in this regime

$$\frac{1}{\phi_k} \frac{d^2 \phi_k}{d\eta^2} = \frac{\ddot{a}}{a}. \quad (88)$$

The solution is therefore given by a growing and a decaying mode. Clearly, to estimate the power spectrum, the growing mode is the most important one:

$$\phi_k(\eta \rightarrow 0) \sim C_k a(\eta). \quad (89)$$

From this last expression, one simply rewrites the power spectrum as a function of C_k :

$$\mathcal{P}_T(k) = \frac{2k^3}{\pi^2} |C_k|^2. \quad (90)$$

B. Results

For the numerical investigations, the solution is first taken as Hankel functions of order 0, given by Eq. (83), in the limit $\eta \rightarrow -\infty$. It is then numerically propagated according to Eq. (5) with a variable step fourth-order Runge-Kutta code. When $\eta \rightarrow 0$, the numerical results are fitted with Eq. (89) and the resulting value of C_k is used to compute the power spectrum. Figure 1 displays, for $\kappa(1 + \epsilon) = 2$, both the numerical results and the analytical calculation in the IR limit obtained with Eq. (75). The excellent agreement illustrates the reliability of the approach. Our code has also been tested with LQC corrections set to zero: the numerical results are also in agreement with the analytically known prediction of standard slow-roll inflation.

A few comments can be made about the general structure of Eq. (8), with the energy and potential terms given by Eq. (10) and Eq. (11). First, it should be noticed that in the limit $\eta \rightarrow -\infty$, the potential is always subdominant when compared with the energy. This makes meaningful the assumption of "asymptotically free" initial states. However, in the remote past, the LQC correction terms are dominant, both in the energy and in the potential. This is why the vacuum structure in the limit $\eta \rightarrow -\infty$ is far from obvious and the intuitive interpretation of the result is very




FIG. 2: Primordial tensor power spectrum as a function of the wavenumber for different values of the slow-roll parameter ϵ .




FIG. 3: Primordial tensor power spectrum as a function of the wavenumber for different values of the LQC parameters λ (left panel) and κ (right panel).

difficult. Furthermore, if $\kappa(1 + \epsilon) > 2$ the potential diverges when $\eta \rightarrow -\infty$. This unusual feature remains harmless as the energy term is, whatever the parameters, much higher than the potential term. When $\eta \rightarrow 0$, *i.e.* at the end of inflation, both the energy and the potential are dominated by the GR terms.

Figures 2, Fig. 3 and Fig. 4 display the primordial tensor power spectrum as a function of the wavenumber for different values of the physical parameters. Our fiducial model is defined by $\epsilon = 0$, $\lambda = 1$, $\kappa = 2$, $H_0 = 0.1$. We remind that ϵ controls the slow-roll whereas λ and κ are LQC parameters defined by Eq. (9) and H_0 is such that $H_0 \equiv \ell_{Pl}/\ell_0$ with $a(\eta) = \ell_0 |\eta|^{-1-\epsilon}$. The main result of this article, which appears in all the figures, can easily be noticed: the spectrum is strongly infrared divergent due to LQC corrections. On the other hand, as expected, the




FIG. 4: Primordial tensor power spectrum as a function of the wavenumber for different values of the parameter H_0 .

UV limit coincides with the GR prediction.

Fig. 2 aims at underlining the general evolution tendency of the spectrum as a function of the slow-roll parameter and the last value ($\epsilon=0.1$) is deliberately chosen above the observationally allowed range [5]. Although it cannot be easily seen on the plot due to the scale, the spectrum exhibits, as it should, a -2ϵ tilt in the UV limit. On Fig. 3, one can notice that, as expected, the higher the value of the LQC parameters, the stronger the deviation from GR. Finally, the case of Fig. 4 is slightly more intricate as H_0 is fundamentally a background parameters which, however, couples to LQC corrections *via* the λ term in the IR regime.

On Fig. 5, the critical wavenumber k_{crit} is displayed as a function of the parameters. We define k_{crit} so that $\mathcal{P}_T(k_{crit}) = 2\lim_{k \rightarrow \infty} \mathcal{P}_T(k)$ (except for a nonvanishing ϵ , in which case the criterion is $\mathcal{P}_T^{(LQG)}(k_{crit}) = 2\mathcal{P}_T^{(RG)}(k_{crit})$), as an indicator of the transition wavenumber between the LQC-dominated ($k < k_{crit}$) and the GR-dominated ($k > k_{crit}$) regimes. Interestingly, it can be concluded from those plots that this critical wavenumber is highly dependent upon any LQC parameter. To some extent, this feature is more observationally relevant than the amplitude of the effect which is anyway huge in the infrared limit. The higher the LQC correction, the higher the critical wavenumber, the smaller the physical scales submitted to LQC corrections, and the easier the observation. Although probably fortuitous, it is worth noticing that the transition scale is nearly proportional to the energy scale of inflation. The dependence upon the first slow-roll parameter is very weak in the allowed range, making the predictions reliable from the viewpoint of a test of LQC.

CONCLUSION

The influence of holonomy corrections during slow-roll inflation was derived in [7, 8]. This article follows the same approach but considers inverse-volume terms (complementing the approach of [35] which was performed in the framework of superinflation). Both analytical results (for a few particular cases) and numerical results (sampling the full parameter space) were obtained. The general behavior is a very substantial deviation from GR in the $k \rightarrow 0$ limit. This deviation affects the amplitude of the power spectrum and, more importantly, both the tilt and the running of the index. There are several ways in which this work should be developed. First, it would be welcome to include simultaneously holonomy and inverse-volume corrections in the same differential equation for primordial gravitons. Then, and more importantly, corrections to the background should also be taken into account. Although several articles are devoted to this point, no fully consistent numerical study of LQC propagation *and* background corrections is yet available. Finally, it would be interesting to investigate LQC corrections to the scalar spectrum [36]. This last




FIG. 5: Value of the critical wavenumber as a function of the inflationary parameters ϵ (top-left) and H_0 (top-right) and as a function of the LQC parameters λ (bottom-left) and κ (bottom-right).

point is probably the most promising one from the observational viewpoint.

-
- [1] C. Rovelli, *Quantum Gravity*, Cambridge, Cambridge University Press, 2004.
 - [2] L. Smolin, arXiv:hep-th/0408048v3.
 - [3] M. Bojowald, Living Rev. Rel. **11**, 4 (2008).
 - [4] A. Linde, Lect. Notes Phys. **738**, 1 (2008).
 - [5] E. Komatsu *et al.*, Astrophys. J. Suppl. Ser. **180**, 330 (2009).
 - [6] S. Tsujikawa, P. Singh, and R. Maartens, Class. Quant. Grav. **21**, 5767 (2004).
 - [7] A. Barrau and J. Grain, Proc. of the 43rd Rencontres de Moriond, arXiv:0805.0356v1[gr-qc].
 - [8] J. Grain and A. Barrau, Phys. Rev. Lett **102**, 081301 (2009).
 - [9] M. Bojowald, P. Singh & A. Skirzewski, Phys. Rev. D **70**, 124022 (2004).
 - [10] P. Singh and K. Vandersloot, Phys. Rev. D **72**, 084004 (2005).

- [11] K. Vandersloot, Phys. Rev. D **71**, 103506 (2005).
- [12] G. Date and G. M. Hossain, Class. Quant. Grav. **21**, 4941 (2004).
- [13] D. Mulryne and N. Nunes, Phys. Rev. D **74**, 083507 (2006).
- [14] T. Thiemann, Class. Quant. Grav. **15**, 839 (1998).
- [15] T. Thiemann, Class. Quant. Grav. **15**, 1281 (1998).
- [16] M. Bojowald, Phys. Rev. D **64**, 084018 (2001).
- [17] M. Bojowald, Living Rev. Rel. **8**, 11 (2005).
- [18] M. Bojowald, Pramana **63**, 765 (2004).
- [19] J. Mielczarek and M. Szydlowski, Phys. Lett. B **657**, 20 (2007); J. Mielczarek, J. Cosmo. Astropart. Phys. 11 (2008) 011.
- [20] M. Bojowald *et al.*, Phys. Rev. D **70**, 043530 (2004).
- [21] J.E. Lidsey, D.J. Mulryne, N.J. Nunes, and Tavakol, Phys. Rev. D **70**, 063521 (2004).
- [22] D.J. Mulryne, N.J. Nunes, R. Tavakol, and J.E. Lidsey, Int. J. Mod. Phys. A **20**, 2347 (2005).
- [23] D.J. Mulryne, R. Tavakol, J.E. Lidsey and G.F.R. Ellis, Phys. Rev. D **71**, 123512 (2005).
- [24] N.J. Nunes, Phys. Rev. D **72**, 103510 (2005).
- [25] E.J. Copeland, D.J. Mulryne, N.J. Nunes, and M. Shaeri, Phys. Rev. D **77**, 023510 (2008).
- [26] M. Bojowald, Phys. rev. Lett. **89**, 261301 (2002).
- [27] M. Bojowald and K. Vandersloot, Phys. Rev. D **67**, 124023 (2003).
- [28] E.J. Copeland, D.J. Mulryne, N.J. Nunes, and M. Shaeri, Phys. Rev. D **79**, 023508 (2009).
- [29] M. Bojowald and G.M. Hossain, Phys. Rev. D **77**, 023508 (2008).
- [30] M. Bojowald, J. Phys. Conf. Ser. **24**, 77 (2005).
- [31] P. Singh, K. Vandersloot, and G.V. Vereshchagin, Phys. Rev. D **74**, 043510 (2006).
- [32] M. Bojowald, Sci. Am. **299N4**, 28 (2008).
- [33] J. Martin, Lect. Notes Phys. **738**, 193 (2008); J. Martin and D. Schwarz, Phys. Rev. D **62**, 103520 (2000); Phys. Rev. D **67** 083512 (2003).
- [34] A. Corichi and P. Singh, Phys. Rev. D **78**, 024034 (2008).
- [35] G. Calcagni and G.M. Hossain, arXiv:0810.4330v3.
- [36] M. Bojowald, G. Hossain, M. Kagan & S. Shankaranarayanan, Phys. Rev. D **79** 043505 (2009).